

Prostat Kanserinde Devam Eden Klinik Çalışmalar Gelecek Perspektif

Dr. Deniz Tural
Bakırköy Dr. Sadi Konuk Eğitim ve Araştırma Hastanesi
Tıbbi Onkoloji

Ders Planı

- ❑ Evre IV kastrasyona Dirençli Prostat Kanserinde Gelecek perspektif
- ❑ Evre IV kastrasyona Duyarlı Prostat Kanserinde Gelecek perspektif
- ❑ Yüksek Riskli Prostat Kanserinde Gelecek perspektif
- ❑ Lokalize Nüks/Biyokimyasal Nükslerde Gelecek Perspektif
- ❑ Sonuç

Evre IV kastrasyona Dirençli Prostat Kanserinde Gelecek perspektif

❑ DNA repair gen mutasyonları


❑ AR mutasyonları

❑ PI3K-PTEN-AKT yolağı

❑ PSMA ekspresyonu

❑ Mikrosatellit instabilite(MSI)

❑ Diğer kombinasyonlar



Cell 2015

Prostat Kanseri Patogenezi

Recent insights into the molecular landscape of advanced PC have identified the following potentially actionable targets:

Molecular alteration	Frequency of expression in advanced PC*
High levels of PSMA expression	 (>80%) ¹⁻⁵
AR pathway mutations/alterations	 (63%–71%) ⁶
PTEN-PI3K-AKT pathway alterations	 (49%) ⁶
Cell cycle (CDK) pathway alterations	 (21%) ⁶
DNA repair pathway alterations	 (19%–23%) ⁶
WNT pathway alterations	 (18%) ⁶
MSI-H, dMMR	 (~3–5%) ^{7,8}

PSMA appears to be the most broadly applicable potential biomarker and actionable target in advanced PC¹⁻⁶

*Each figure represents 10% of patients with advanced PC.

1. Hope TA, et al. *J Nucl Med.* 2017;58(12):1956–1961; 2. Hupe MC, et al. *Front Oncol.* 2018;8:623; 3. Pomykala KL, et al. *J Nucl Med.* 2020;61(3):405–411; 4. Minner S, et al. *Prostate.* 2011;71(3):281–288; 5. Bostwick DG, et al. *Cancer.* 1998;82(11):2256–2261; 6. Robinson D, et al. *Cell.* 2015;161(5):1215–1228; 7. Abida W, et al. *JAMA Oncol.* 2019; 5(4):471–478; 8. Lindh C, et al. *APMIS.* 2019; 127(8):554–560.

AKT, protein kinase B; AR, androgen receptor; CDK, cyclin-dependent kinase; PC, prostate cancer; PI3K, phosphoinositide 3-kinase; PSMA, prostate-specific membrane antigen; PTEN, phosphatase and tensin homolog; WNT, wingless int-1.

Evre IV Kastrasyona Dirençli Prostat Kanseri

Ipatasertib and the PI3K-AKT Pathway

PI3K/AKT pathway biomarkers analysis from the Phase III IPATential150 trial of ipatasertib plus abiraterone in metastatic castration-resistant prostate cancer


Johann de Bono,¹ Christopher Sweeney,² Sergio Bracarda,³ Cora N. Sternberg,⁴ Kim N. Chi,⁵ David Olmos,⁶ Shahneen Sandhu,⁷ Christophe Massard,⁸ Nobuaki Matsubara,⁹ Josep Garcia,¹⁰ Małgorzata Nowicka,¹⁰ Matthew Wongchenko,¹¹ Zhen Shi¹¹

1. The Institute of Cancer Research and the Royal Marsden Hospital, London, UK; 2. Dana-Farber Cancer Institute, Boston, MA, USA; 3. Azienda Ospedaliera S. Maria, Terni, Italy; 4. Englewood Institute for Precision Medicine, Dana-Farber Cancer Institute, Boston, MA, USA; 5. BC Cancer Agency, Vancouver, Canada; 6. Spanish National Cancer Research Center (CNIO), Madrid, Spain; 7. Peter MacCallum Cancer Centre, Melbourne, Australia; 8. Institut Gustave Roussy, Villejuif, France; 9. National Cancer Center Hospital East, Chiba, Japan; 10. F. Hoffmann-La Roche, Ltd., Basel, Switzerland; 11. Genentech, South San Francisco, CA, USA.

Presented at: Genitourinary Cancers Symposium | Boston, MA | December 10-12, 2021 | 40021

Presented by: Johann de Bono, MD
<https://bit.ly/3b1y2P>


Presented By Johann De Bono at 2021 Genitourinary Cancers Symposium



J.S. Brown, U. Banerji / Pharmacology & Therapeutics 172 (2017) 101–115

Evre IV Kastrasyona Dirençli Prostat Kanseri

IPATential150 Study Design and Co-primary Endpoint Outcomes



Investigator-assessed rPFS in the PTEN-loss (by IHC) population

	Pbo + abi n = 261	Ipat + abi n = 260
Patients with event, n (%)	154 (59)	124 (48)
1-Year event-free rate (95% CI), %	63.3 (57.3, 69.3)	64.4 (58.3, 70.5)
Stratified HR (95% CI)	0.77 (0.61, 0.98); $P = 0.0335^c$	


Investigator-assessed rPFS in the ITT population

	Pbo + abi n = 554	Ipat + abi n = 547
Patients with event, n (%)	306 (55)	252 (46)
1-Year event-free rate (95% CI), %	63.0 (58.9, 67.1)	65.3 (61.1, 69.5)
Stratified HR (95% CI)	0.84 (0.71, 0.99); $P = 0.0431^d$	

HSPC, hormone-sensitive prostate cancer; IHC, immunohistochemistry; NGS, next-generation sequencing; PCWG3, Prostate Cancer Working Group 3; R, randomized.
^a PTEN loss status was defined as a minimum of 50% of the specimen's tumor area with no detectable PTEN staining (by Ventana PTEN (SP218) Investigational IHC assay using SP218 antibody). ^b Abiraterone s (1000 mg qd) plus prednisone/prednisolone (5 mg bid). ^c Statistically significant at $\alpha = 0.05$ level. ^d Did not meet statistical significance at $\alpha = 0.01$ level.

Evre IV Kastrasyona Dirençli Prostat Kanseri

Improvement in rPFS by NGS Population



PTEN WT by NGS (among PTEN NGS evaluable)	Pbo + abi n = 155	Ipat + abi n = 155
rPFS, median (95% CI), mo	16.6 (13.7, 24.9)	20.9 (16.4, NE)
Stratified HR (95% CI)	0.85 (0.62, 1.18)	

PTEN WT + PTEN loss by NGS (among PTEN NGS evaluable)	Pbo + abi n = 258	Ipat + abi n = 260
rPFS, median (95% CI), mo	16.5 (13.7, 19.0)	19.3 (16.7, 24.7)
Stratified HR (95% CI)	0.77 (0.60, 0.99)	

¹OS data were immature; however, the investigators noted an early trend showing survival favoring the ipatasertib arm in the PTEN-loss group (HR, 0.91) and ITT population (HR, 0.93)

L. de Bono JS, Bracarda S, Sternberg CN, et al. IPATential150: efficacy and safety from the phase III study of ipatasertib plus abiraterone vs placebo plus abiraterone in metastatic castration-resistant prostate cancer. Presented at: 2020 ESMO Virtual Congress; September 19-21, 2020; Virtual. Abstract LBA4.

Presented By Johann De Bono at 2021 Genitourinary Cancers Symposium

Evre IV Kastrasyona Duyarlı Prostat Kanseri


FUTURE PERSPECTIVES

Precision medicine in mHSPC

	Intervention	Control Arm	Column C
TALAPRO-3 NCT04821622	Talazoparib + Enzalutamide	Enzalutamide	DDR gene mutated
CAPItello-281 NCT04493853	Capivasertib + Abi	Abi	PTEN deficiency

Evre IV Kastrasyona Dirençli Prostat Kanseri

RATIONALE FOR TREATMENT IN PROSTATE CANCER




Cell cycle disruption

Evre IV Kastrasyona Dirençli Prostat Kanseri


CYCLONE 2

Phase II/III, RCT, placebo controlled



Evre IV Kastrasyona Dirençli Prostat Kanseri

> 1 in 10 Men with mCRPC have Germline Mutations



- Inherited germline DDR mutations
 - mCRPC: 11.8% (82/692)
 - Localized disease: 4.6% (23/499)

Presumed pathogenic germline mutations in metastatic cases (N = 692)		
Gene	No. of Mutations	% of Men
<i>BRCA2</i>	37	5.35
<i>ATM</i>	11	1.59
<i>CHEK2*</i>	10	1.87
<i>BRCA1</i>	6	0.87

*n = 534; data censored for metastatic cases with inadequate sequencing



Using germline testing alone
~50% of patients with a BRCA1/2, or ATM mutation will be missed

Evre IV Kastrasyona Dirençli Prostat Kanseri

Patients with HRRm including *BRCA2*m are more likely to have poor outcomes on standard of care therapies¹⁻³

Patients with **germline HRRm** including *BRCA2*m are more likely to have **poor outcomes** on standard of care therapies^{1,2}


Poor responses to standard therapy also seen for **tumour HRRm³**



1. Castro E, et al. *J Clin Oncol*. 2019;6:490–503; 2. Annala M, et al. *Eur Urol*. 2017;72:34–42; 3. Annala M, et al. *Cancer Discovery*. 2018;doi:10.1158/2159-8290.CD-17-0937

Evre IV Kastrasyona Dirençli Prostat Kanseri

PARP Inhibitor Trials and approvals in mCRPC



HRR, high level results; HRRm, homologous recombination repair; mCRPC, metastatic castration-resistant prostate cancer; PARP, poly(ADP-ribose) polymerase; P, phase

1. de Bono J et al. *NEJM* 2020;382:2091-102; 2. AstraZeneca press release, 7 August 2019; 3. <https://www.fda.gov/drugs/drug-approvals-and-databases/fda-approves-olaparib-hcr-gene-mutated-metastatic-castration-resistant-prostate-cancer>; 4. Mateo J et al. *NEJM* 2015;

575:1697-706; 5. Mateo J et al. *J Clin Oncol* 2019;37:Abstr 5005; 6. <https://clinicaltrials.gov/ct2/show/NCT02292334>

7. <https://clinicaltrials.gov/ct2/show/NCT02975954>; 8. <https://www.fda.gov/drugs/fda-grants-accelerated-approval-rucaparib-hcr-mutated-metastatic-castration-resistant-prostate>


9. <https://clinicaltrials.gov/ct2/show/NCT03148793>; 10. <https://clinicaltrials.gov/ct2/show/NCT02804428>; 11. Lynparza [olaparib] Canadian Product Monograph.

Evre IV Kastrasyona Dirençli Prostat Kanseri


PROfound Secondary Endpoint: Significant Improvement in OS in mCRPC with BRCA1/2 or ATM Mutations (Cohort A)

31% Reduction in Risk Of Death with Olaparib vs. Physician's Choice

Cohort A



Cohort A with adjustment for crossover*




Median follow-up duration for censored patients: olaparib, 21.9 months; control, 21.0 months. *Re-censored; conducted using rank-preserving structural failure time model (RPSTF) to demonstrate the impact on OS of crossover of patients from the control arm to receive olaparib as a first subsequent anticancer therapy. CI, confidence interval; HR, hazard ratio; OS, overall survival. 1. Hussain M, et al. *NEJM* 2020;Online: doi10.1056/NEJMoa2022485

Evre IV Kastrasyona Dirençli Prostat Kanseri

PROpel

Randomized, double-blind, placebo-controlled Phase III trial



Baseline demographics:
HRRm status

HRRm status*		
HRRm	111 (27.8)	115 (29.0)
Non-HRRm	279 (69.9)	273 (68.8)
HRRm unknown	9 (2.3)	9 (2.3)



Evre IV Kastrasyona Dirençli Prostat Kanseri


PROpel

Primary endpoint

rPFS by investigator assessment



rPFS by blinded independent central review^




- 34% risk reduction for progression or death with olaparib + abiraterone (HR 0.66; 95% CI 0.54–0.81; P<0.0001)

Evre IV Kastrasyona Dirençli Prostat Kanseri

MAGNITUDE



Randomized, double-blind, placebo-controlled Phase III trial



Evre IV Kastrasyona Dirençli Prostat Kanseri

MAGNITUDE


Primary endpoint: rPFS by central review



- 47% improvement in rPFS in patients with *BRCA1/2* alterations (HR 0.53; 95% CI 0.36–0.79; $P=0.0014$)
- 27% improvement in rPFS across all HRR BM+ patients (HR 0.73; 95% CI 0.56–0.96; $P=0.0217$)

Evre IV Kastrasyona Dirençli Prostat Kanseri

TALAPRO-2: Phase III Trial Design – Talazoparib + Enzalutamide



www.clinicaltrials.gov: (NCT03395197)

Agarwal,..Fizazi. TALAPRO-2 Phase III study design, Future Oncology 2022.


Evre IV Kastrasyona Dirençli Prostat Kanseri



- *TALZENNA[®] first PARP inhibitor to demonstrate clinical benefit in combination with XTANDI[®] in metastatic castration-resistant prostate cancer (mCRPC)*
- *Study achieves primary endpoint of radiographic progression-free survival*
- *Robust, highly consistent efficacy demonstrated in mCRPC both with or without homologous recombination repair gene mutations*

Evre IV Kastrasyona Dirençli Prostat Kanseri

Ongoing trials investigating PARPi in advanced PC



Please see slide notes for references

*As a result of the data from PROfound, olaparib monotherapy was approved for treatment of mCRPC in patients with HRRm (FDA approval) or for patients with mutations in only BRCA1/2 (EMA approval) after progression on a NHA^{12,13}

[†]As a result of the data from TRITON2, rucaparib monotherapy was approved by the FDA only for the treatment of mCRPC in patients with a BRCA1/2m who have disease progression after treatment with prior AR-directed therapy and prior taxane¹⁴


Abi=abiraterone; BCR=biochemical recurrence; enza=enzalutamide; FDA=US Food and Drug Administration; HRRm=homologous recombination repair mutation; mCRPC=metastatic castration-resistant prostate cancer; met=metastasis;

mono=monotherapy; mHSPC=metastatic hormone-sensitive prostate cancer; NHA=new hormonal agent; nmCRPC=non-metastatic castration-resistant prostate cancer; P2=Phase II; P3=Phase III.


Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA ekspresyonu

EAU22 | AMSTERDAM
1-4 July 2022

Synergism: Holy Grail



Sartor et al., NEJM 2021



<https://biology-forums.com/index.php?action=gallery;sa=view;id=28366>

$$1+3 = 4$$

$$1+3 > 4$$

Evre IV Kastrasyona Dirençli Prostat Kanseri

PSMA ekspresyonu


EAU22 | AMSTERDAM
1-4 July 2022

Candidates

- Checkpoint Inhibitors
- PARP Inhibitors
- ARDT
- Chemotherapy
- SBRT
- Others?


Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA ekspresyonu

PSMA-Lutetium Radionuclide Therapy and ImmuNotherapy for Prostate CancEr (PRINCE) Trial Schema



Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA ekspresyonu

PRINCE: PSA Response Rate (Primary Endpoint)



Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA ekspresyonu


EAU22 | AMSTERDAM
1-4 July 2022

PARP Inhibitors

Combination with Olaparib

LuPARP Trial

Phase 1 trial of ^{177}Lu -PSMA-617 therapy and **Olaparib (PARPi)**




Evre IV Kastrasyona Dirençli Prostat Kanseri

PSMA ekspresyonu

EAU22 | AMSTERDAM
1-4 July 2022

ARDT

ENZA-p Trial




Emmett et al., BJUI 2021

PSMAaddition

PSMAaddition: a randomized, phase 3 study of ^{177}Lu -PSMA-617 in patients with untreated or minimally treated mHSPC

- PSMAaddition aims to assess the efficacy and safety of ^{177}Lu -PSMA-617 RLT plus SoC^{*} vs SoC in men with untreated/minimally treated mHSPC (NCT04720157)¹




Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA ekspresyonu

EAU22 | AMSTERDAM
1-4 July 2022

Chemotherapy

UpfrontPSMA (NCT04343885)



Evre IV Kastrasyona Dirençli Prostat Kanseri

PSMA ekspresyonu

EAU22 | AMSTERDAM
1-4 July 2022

SBRT

Combining SBRT and PSMA RLT

A Study of Stereotactic Body Radiotherapy and ¹⁷⁷Lu-PSMA-617 for the Treatment of Prostate Cancer

ClinicalTrials.gov Identifier: NCT05079698

Recruitment Status ⓘ : Recruiting

First Posted ⓘ : October 15, 2021

Last Update Posted ⓘ : June 6, 2022


See [Contacts and Locations](#)

Sponsor:

Memorial Sloan Kettering Cancer Center

Collaborator:

Novartis Pharmaceuticals



Rationale of „adding“ SBRT in patients with few, disease dominant but PSMA-negative or PSMA-low uptake lesions very appealing..

Kuo et al., SNMMI 2022


Evre IV Kastrasyona Dirençli Prostat Kanseri

PSMA ekspresyonu

EAU22 | AMSTERDAM
1-4 July 2022

Others

Current Lu-PSMA combination trials




Evre IV Kastrasyona Dirençli Prostat Kanseri

Mikrosatellit instabilite

Immune checkpoint blockade

**Limited activity of PD-1/PD-L1 inhibitors in CRPC?
(mutations are rare)**



Alexandrov et al, Nature 2013


for re-use.



Evre IV Kastrasyona Dirençli Prostat Kanseri

İmmüne checkpoint İnhibitörleri

IMPACT: Phase III Trial of Sipuleucel-T




Primary endpoint: Overall survival

Secondary endpoint: Time to objective disease progression

†64% of patients in the control group, following progression, crossed over to receive autologous immunotherapy made from cryopreserved cells.

Evre IV Kastrasyona Dirençli Prostat Kanseri İmmune Checkpoint İnhibitörleri

Sipuleucel-T: Impact Phase 3 trial A first « negative-positive » trial



Evre IV Kastrasyona Dirençli Prostat Kanseri İmmüne Checkpoint İnhibitörleri



The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

PD-1 Blockade in Tumors with Mismatch-Repair Deficiency

D.T. Le, J.N. Uram, H. Wang, B.R. Bartlett, H. Kemberling, A.D. Eyring, A.D. Skora, B.S. Luber, N.S. Azad, D. Laheru, B. Biedrzycki, R.C. Donehower, A. Zaheer, G.A. Fisher, T.S. Crocenzi, J.J. Lee, S.M. Duffy, R.M. Goldberg, A. de la Chapelle, M. Koshiji, F. Bhajee, T. Huebner, R.H. Hruban, L.D. Wood, N. Cuka, D.M. Pardoll, N. Papadopoulos, K.W. Kinzler, S. Zhou, T.C. Cornish, J.M. Taube, R.A. Anders, J.R. Eshleman, B. Vogelstein, and L.A. Diaz, Jr.


2% of prostate cancers with MMR mutations




Evre IV Kastrasyona Dirençli Prostat Kanseri İmmüne Checkpoint İnhibitörleri

Inactivation of *CDK12* Delineates a Distinct Immunogenic Class of Advanced Prostate Cancer

Yi-Mi Wu,^{1,2,20} Marcin Cieřlik,^{1,2,20} Robert J. Lonigro,¹ Pankaj Vats,¹ Melissa A. Reimers,³ Xuhong Cao,¹ Yu Ning,¹ Lisha Wang,¹ Lakshmi P. Kunju,^{1,2,4} Navonil de Sarkar,⁵ Elisabeth I. Heath,^{6,7} Jonathan Chou,⁸ Felix Y. Feng,^{8,9,10,11} Peter S. Nelson,^{5,12,13} Johann S. de Bono,^{14,15} Weiping Zou,^{1,2,16} Bruce Montgomery,^{12,17} Ajjai Alva,^{1,3} PCF/SU2C International Prostate Cancer Dream Team, Dan R. Robinson,^{1,2,*} and Arul M. Chinnaiyan^{1,2,4,18,19,21,*}



- 6.9% [4.6%, 10.2%] of mCRPC
- Focal tandem duplications.¹





Evre IV Kastrasyona Dirençli Prostat Kanseri İmmune Checkpoint İnhibitörleri

Pilot Clinical Study to Determine CDK12 Mutant Prostate Cancer
Response to Checkpoint Inhibitor Immunotherapy

11 pts -5 tt anti PD1

- 1 pt excluded
- 2 pts +++PSA decline



Evre IV Kastrasyona Dirençli Prostat Kanseri Immune Checkpoint İnhibitörleri

Metastatic Castration-resistant Prostate Cancer	Drug: Atezolizumab	
--	--------------------	--

Detailed Description:

This is a multi-center, open label Phase II study of patients with metastatic castration resistant prostate cancer (mCRPC) who will be treated with abemaciclib and atezolizumab alone or in combination.

The U.S. Food and Drug Administration (FDA) has not approved abemaciclib or atezolizumab alone or in combination for use in prostate cancer. Abemaciclib is an orally administered molecularly targeted chemotherapy drug called a cyclin-dependent kinase inhibitor, which acts to block the ability of cancer cells to divide and thus prevents tumors from growing. In the laboratory setting, this drug is effective in prostate cancer models that have become resistant to standard hormonal treatments, and this drug is currently being studied for its effectiveness in prostate cancer in other clinical trials. Atezolizumab is an intravenously administered drug called an immune checkpoint inhibitor, which acts to activate the immune system to kill cancer cells. Atezolizumab is ineffective on its own in most patients with prostate cancer, but is being tested in combination with other drugs for prostate cancer in other clinical trials. Multiple research groups have demonstrated in laboratory model systems that abemaciclib can may make immune checkpoint inhibitors more effective.

The research study procedures include screening for eligibility and study treatment including evaluations and follow up visits.

The study design divides study participants into two separate cohorts. The first cohort is a set of subjects whose tumors are not known to have mutations in the CDK12 gene (the "biomarker unselected cohort") - either because tumor tissue never underwent genetic profiling, or because genetic profiling was performed but did not demonstrate a mutation in the CDK12 gene. In this "biomarker unselected cohort," this study will be testing whether abemaciclib alone or in combination with atezolizumab is an effective treatment strategy.

The second cohort of participants is a set of subjects whose tumors are known to have mutations in the CDK12 gene based on genetic profiling of the tumor that occurred prior to enrollment on this study. Prior studies suggest that cancers with mutations in the CDK12 gene can shrink in response to immune checkpoint inhibitors. This study will be testing in study participants whose tumors are known to have mutations in CDK12 whether atezolizumab alone or in combination with abemaciclib is an effective treatment strategy.

In addition, the trial is testing the safety of the combination of the two drugs in both cohorts.


Participants will receive study treatment for as long as they do not have serious side effects and their disease does not get worse. Participants will be followed after completion of study treatment for up to 24 months

It is expected that about 75 people will take part in this research study.


Eli Lilly and Company is supporting this research study by providing funding for research and the study drug abemaciclib. Genentech, Inc. is supporting the study by providing the study drug atezolizumab.

Evre IV Kastrasyona Dirençli Prostat Kanseri İmmun Checkpoint İnhibitörleri

Keynote 365 – Cohort B: Pembrolizumab + Enzalutamide




Conter et al., ASCO 2020




Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.

Evre IV Kastrasyona Dirençli Prostat Kanseri İmmun Checkpoint İnhibitörleri

Keynote 365 – Cohort B: Pembrolizumab + Docetaxel



Sridhar et al., ASCO 2020






Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.



Evre IV Kastrasyona Dirençli Prostat Kanseri İmmun Checkpoint İnhibitörleri


COSMIC-021 phase 1b: Atezolizumab + Cabozantinib




Evre IV Kastrasyona Dirençli Prostat Kanseri İmmün Checkpoint İnhibitörleri

CHECKMATE 9KD: Nivolumab + Rucaparib

Median r PFS



Median OS



Evre IV Kastrasyona Dirençli Prostat Kanseri

İmmüne Checkpoint İnhibitörleri


Immune Checkpoint Inhibitors in mCRPC

Therapy	Disease State	Disease Response
Pembrolizumab monotherapy ^a	Post-chemotherapy	ORR 9% PSA RR 14%
Pembrolizumab + enzalutamide ^b	Pre-chemotherapy progressing on enzalutamide	ORR 12% PSA RR 14%
Atezolizumab + enzalutamide ^c	Pre- and post-chemotherapy, s/p abiraterone	ORR 14% PSA RR 26%
Atezolizumab + cabozantinib ^d	Pre-chemotherapy s/p enzalutamide or abiraterone	ORR 34% PSA RR 29%

^aJCO 2020: 38(5) 395-405. ^bPresented at the 2021 ASCO Annual Meeting – Virtual; June 4-8, 2021. ^cSweeney C. AACR 2020. IMbassador250. ^dAgarwal ASCO 2020. COSMIC-021

Evre IV Kastrasyona Dirençli Prostat Kanseri İmmüne Checkpoint İnhibitörleri

CONTACT-02: Phase III Trial Schema



Stratification

- Liver metastasis (yes, no)
- Prior docetaxel treatment for mCSPC (yes, no)
- Disease stage for which the first NHT was given (mCSPC, M0 CRPC, mCRPC)

*Second NHT must differ from previous NHT taken


[†]Tumor assessment (RECIST v1.1) every 9 weeks for the first 28 weeks then every 12 weeks thereafter

[‡]Patients may be treated beyond progression if there is a clinical benefit in the opinion of the investigator

Evre IV Kastrasyona Dirençli Prostat Kanseri

PSMA BİTE

Targeting PSMA: PSMA BiTE



BiTE molecules engage a patient's own T cells to attack and eradicate cancer cells

— T-cell activation induces transient cytokine release and tumor killing

1. Baeuerle PA, et al. *Cancer Res.* 2009; 2. Klinger M, et al. *Immun Rev.* 2016; 3. Bargou R, et al. *Science.* 2008;
4. Stieglmaier J, et al. *Expert Opin Biol Ther.* 2015;15(8):1093-9.

Evre IV Kastrasyona Dirençli Prostat Kanseri PSMA BİTE



AMG 160 monotherapy: Results

PSA/CTC Responses (n = 13–35)

Response	All, n (%)
PSA response, confirmed*	8 (27.6)
PSA response, unconfirmed†	4 (11.4)
CTC0 response‡	3 (23.1)

RECIST Responses (n = 15)

Response	All, n (%)
Partial response, confirmed	2§ (13.3)
Partial response, unconfirmed	1§ (6.7)
Stable disease	8 (53.3)



CTC = circulating tumor cell; DLT = dose-limiting toxicities; NE = not evaluable; PSA = prostate-specific antigen; PR = partial response; Q2W = every 2 weeks; RECIST = Response Evaluation Criteria in Solid Tumors; SD = stable disease; (u) = unconfirmed


Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.



Evre IV Kastrasyona Dirençli Prostat Kanseri


PSMA BİTE

HPN424: “Trispecific T-cell Activating Constructs” (TriTACs)



Anti-Tumor Activity


Treatment duration of >24 weeks in 20% of patients, Confirmed RECIST PR, PSA declines and CTC reductions observed across cohorts



Patient 057 Target Lesion Scans




Patient 054 Target Lesion Scans



Evre IV Kastrasyona Dirençli Prostat Kanseri

CAR T Cells



Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.

Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar




A phase 3 trial with a 2x2 factorial design in men with *de novo* metastatic castration-sensitive prostate cancer (mCSPC): Overall survival with abiraterone acetate plus prednisone in PEACE-1

Karim Fizazi, Joan Carles, Stéphanie Foulon, Guilhem Roubaud, Ray McDermott, Aude Fléchon, Bertrand Tombal, Stéphane Supiot, Dominik Berthold, Philippe Ronchin, Gabriel Kacsó, Gwenaëlle Gravis, Fabio Calabro, Jean-François Berdah, Ali Hasbini, Marlon Silva, Antoine Thiery-Vuillemin, Igor Latorzeff, Isabelle Rieger, Alberto Bossi

Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar


PEACE-1: Radiographic PFS (rPFS) by Metastatic Burden



Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar

ARASENS Primary Endpoint*: Overall Survival

Darolutamide significantly reduced the risk of death by 32.5%





*Primary analysis occurred after 533 deaths (darolutamide, 229; placebo, 304). CI, confidence interval; NE, not estimable.

Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar

ARASENS: ADT + docetaxel +/- darolutamide Overall Survival By Metastatic Stage at Initial Diagnosis

22



Evre IV Kastrasyona Duyarlı Prostat Kanseri İkili Kombinasyonlar

Doublet Therapy Trials in mHSPC: Results

Trial	Patients Enrolled	Intervention Arm	Control Arm	Prior/ Concurrent Docetaxel	Median Follow-up (mo)	Median OS in Interventional Arm (mo)	Median OS in Control Arm (mo)
CHAARTED¹	790	ADT + Docetaxel		Not allowed	53.7	57.6	47.2
STAMPEDE²	2962	ADT + Docetaxel		Not allowed	78.2	59.1	43.1
LATITUDE³	1199	ADT + Abiraterone + Prednisone		Not allowed	51.8	53.3	36.5
STAMPEDE⁴	1917	ADT + Abiraterone + Prednisone		Not allowed	40	HR, 0.61 (P <0.001)	
ENZAMET⁵	1125	ADT+Enzalutamide	ADT + Placebo	Allowed	68	NR	73.2
ARCHES⁶	1150	ADT+Enzalutamide	ADT + Placebo	Allowed	44.6	HR, 0.66 (P <.0001) (median not reached in either group)	
TITAN⁷	1052	ADT + Apalutamide	ADT + Placebo	Allowed	44	NR	52.2

1.Kyriakopoulos et al. J Clin Oncol 2018; 2.Clarke et al. Ann Oncol 2019; 3-Fizazi et al. Lancet Oncol 2019; 4-James et al. N Engl J Med 2017; 5-Davis et al. ASCO Annual Meeting 2022; 6-Armstrong et al. J Clin Oncol 2022; 7-Chi et al. J Clin Oncol 2021

Evre IV Kastrasyona Duyarlı Prostat Kanseri

Tripler kombinasyonlar


Trials with Triplet Therapy in mHSPC: Results

Trial	Patients Enrolled	Intervention Arm	Control Arm	% De-novo	%High Volume	Median Follow-up (mo)	Median OS in Interventional Arm (mo)	Median OS in Control Arm (mo)
ARASENS ¹	1306	ADT + Docetaxel + Darolutamide	ADT + Docetaxel + Placebo	86		43.7	NR	48.9
PEACE-1 ²	710	ADT + Docetaxel + Abiraterone	ADT + Docetaxel	100	64	45.6	NR	52.8
PEACE-1 Subgroup analysis : De-novo High volume of disease →							61	42

1.Smith et al. N Engl J Med 2022; 2.Fizazi et al. Lancet 2022.

Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar

Phase III Trial: Triplets (ARAT+ Docetaxel + ADT) vs. Docetaxel + ADT




ECOG PS, Eastern Cooperative Oncology Group performance status; ART, Androgen receptor targeted therapy; NA, North America; PSA, prostate-specific antigen; OS, Overall survival; rPFS, radiographic progression-free survival.

Evre IV Kastrasyona Duyarlı Prostat Kanseri Üçlü Kombinasyonlar

This trial has not been done yet:

Triplet (Docetaxel + ARAT + ADT) versus ARAT + ADT




ECOG PS, Eastern Cooperative Oncology Group performance status; ART, Androgen receptor targeted therapy; NA, North America; PSA, prostate-specific antigen; OS, Overall survival; rPFS, radiographic progression-free survival.

Lokalize Yüksek Riskli Prostat Kanseri

Localized High-Risk Prostate Cancer

Patterns of Presentation




Increased Mortality in High-Risk Prostate Cancer

High-Risk Feature	15-Year PCSM
PSA > 20 ng/mL	22%
Gleason 8-10	34%
cT3	38%
High-risk Disease*	19%

Lokalizasyon Yüksek Riskli Prostat Kanseri


Metastasis-Free Survival with the Addition of Abiraterone Acetate and Prednisolone with or without Enzalutamide to ADT for High-Risk M0 Prostate Cancer



Lokalizasyon Yüksek Riskli Prostat Kanseri

Overall Survival with the Addition of Abiraterone Acetate and Prednisolone with or without Enzalutamide to ADT for High-Risk M0 Prostate Cancer


Overall survival



Definitif RT+ADT+ Yeni Nesil AR-yolağı İnhibitörleri

Apalutamide

ATLAS: Apalutamide in high-risk, localized or locally advanced PC patients receiving primary RT




Enzalutamide

ENZARAD: Enzalutamide in ADT with RT for high-risk, clinically localized PC



- Primary endpoint is OS
- Secondary endpoints include CSS, PSA PFS, clinical PFS, time to subsequent hormonal therapy, time to castration-resistant disease (PCWG2 criteria), MFS, AEs and HRQOL

SCHEMA




NRG GU009

Abiraterone + Apalutamide

Clinical Trial Information:
NCT02531516; NCT02446444; NCT04513717

Neoadjuvan Tedavi Gerekçesi

Treatment Paradigms for High-Risk Prostate Cancer



ADT=Androgen deprivation therapy.

• Neoadjuvant Treatment

- Standard of care for breast, rectal, bladder and other cancers given improved long-term survival
- Down-stage local disease, which may facilitate surgical resection
- Reduce or delay post-surgery treatment
- Provide an *in vivo* assessment of response to treatment

Pertrelli et al, Eur Urology, 2014
Berger et al, JCO, 2005
Mass et al, Lancet Oncology, 2010
Cortazar et al, Lancet Oncology, 2014
McKay et al, Drugs, 2012

Neoadjuvan Yeni Nesil AR-yolağı İnhibötörleri

Pathologic Responses from Potent Neoadjuvant Therapy

Conducted a series of neoadjuvant trials over the last 10 years


Phase 2 biomarker integrated trials evaluating potent neoadjuvant hormone therapy

	NeoAbi (n=58)	NeoEnza (n=40)	NeoAbiEnza (n=75)
Arms	12wAbi vs. 24wAbi	Enza vs. EDL	EL vs. APEL
CR	4% vs. 10%	0% vs. 4%	8 vs. 12%
MRD*	0% vs. 14%	0% vs. 13%	11% vs. 18%
CR + MRD	4% vs. 24%	0% vs. 17%	19% vs. 30%

Abi=Abiraterone; Enza=Enzalutamide; EDL=Enzalutamide, dutasteride, leuprolide; APEL=Abiraterone, prednisone, enzalutamide, leuprolide; EL=Enzalutamide, leuprolide; CR=Complete response; MRD=Minimum residual disease. *Defined as residual tumor with largest cross section dimension ≤ 0.3 -0.5 cm.

Taplin et al, JCO, 2014
Montgomery et al, CCR, 2016
McKay et al, JCO, 2019

Neoadjuvan Yeni Nesil AR-yolağı İnhibötörleri



Neoadjuvan Yeni Nesil AR-yolağı İnhibötörleri

Pathologic Outcomes

		APL (n=59)	APAL (n=55)
Pathologic Response	pCR	6 (10%)	7 (13%)
	MRD (≤ 5 mm)*	6 (10%)	5 (9%)
	pCR or MRD	12 (20%)	12 (22%)
ypT stage, n (%)	T0	6 (10%)	7 (13%)
	T2	19 (32%)	21 (38%)
	T3	34 (58%)	27 (49%)
pN1, n (%)		10 (17%)	4 (7%)
Positive surgical margins, n (%)		7 (12%)	4 (7%)
Seminal vesicle invasion, n (%)		16 (27%)	15 (27%)
Percent cellularity, % (range)**		5% (0-50%)	5% (0-80%)
RCB (cm ³)		0.07 (0-6.8)	0.02 (0-7.8)

APAL=Abiraterone, prednisone, apalutamide, leuprolide; APL=Abiraterone, prednisone, leuprolide; pCR=Pathologic complete response; MRD=Minimum residual disease; RCB=Residual cancer burden. *Minimum residual disease was defined as residual tumor in the radical prostatectomy specimen measuring ≤ 5 mm. **Residual cancer burden calculated as tumor volume (cm³) x percent cellularity.

Neoadjuvan Yeni Nesil AR-yolağı İnhibitörleri

Neoadjuvant Novel Hormonal Therapy Followed by Prostatectomy versus Up-Front Prostatectomy for High-Risk Prostate Cancer: A Comparative Analysis

Praful Ravi,¹ Lucia Kwak,¹ Wanling Xie,¹ Kaitlin Kelleher,¹ Andres M. Acosta,¹ Rana R. McKay,² Adam S. Kibel,¹ and Mary-ellen Taplin^{1*}

¹Lank Center for Genitourinary Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts

²Department of Medicine, Division of Hematology-Oncology, University of California, San Diego, La Jolla, California

Purpose: We sought to compare outcomes between neoadjuvant therapy with a novel hormonal agent (NHA) prior to radical prostatectomy (neo-RP) and up-front radical prostatectomy (RP) in patients with high-risk prostate cancer (HRPC).

Materials and Methods: HRPC patients treated on 3 trials of neoadjuvant NHA followed by RP formed the neo-RP cohort (112). The RP group (259) comprised an observational cohort of HRPC patients undergoing RP without neoadjuvant therapy between 2010–2016 at our institution who met key eligibility criteria for the neoadjuvant trials (ie ≥ 3 positive biopsy cores and Gleason $\geq 4+3=7$). Inverse probability of treatment weighting (IPTW) was used to minimize potential confounding factors when estimating treatment effects. The primary outcomes were time to biochemical recurrence (BCR) and metastasis-free survival (MFS).

Results: Before IPTW, the neo-RP cohort had higher rates of Gleason 9-10 cancer (46% vs 24%), cT3 disease (22% vs 5%), and PSA ≥ 20 ng/ml (14% vs 7%); after IPTW, the 2 cohorts were balanced. Overall, after IPTW, time to BCR (HR=0.25 [95% CI 0.18–0.37]) and MFS (HR=0.26 [0.15–0.46]) were significantly longer in the neo-RP compared to the RP cohort. Rates of adjuvant (7% vs 24%) and salvage therapy (34% vs 46%) were lower in the neo-RP cohort.

Conclusions: Neoadjuvant therapy with an NHA prior to RP was associated with longer time to BCR and superior MFS compared to up-front RP in men with HRPC. These findings are hypothesis-generating but suggest benefit with neoadjuvant therapy with an NHA in HRPC, an approach which is currently being studied in the phase 3 PROTEUS trial (NCT03767244).

Abbreviations and Acronyms

ADT = androgen deprivation therapy

BCR = biochemical recurrence

HRPC = high-risk prostate cancer

IPTW = inverse probability of treatment weighting

MFS = metastasis-free survival

MRD = minimal residual disease

NHA = novel hormonal agent

OS = overall survival

pCR = pathological complete response

RP = radical prostatectomy

Key Words: neoadjuvant therapy, prostatic neoplasms, general surgery

Neoadjuvan Yeni Nesil AR-yolağı İnhibitörleri




Figure 1. Time to BCR in the neo-RP and RP cohorts after IPTW. The number of participants at risk are based on unweighted population.





Figure 2. MFS in the neo-RP and RP cohorts after IPTW. The number of participants at risk are based on unweighted population.

Devam Eden Çalışmalar

Proteus

Randomized, Double-blind, Placebo-Controlled, Phase 3 Study of Apalutamide in Subjects with High-risk, Localized or Locally Advanced Prostate Cancer Who are Candidates for Radical Prostatectomy




Neoadjuvan Yeni Nesil AR-yolağı İnhibitörleri

Tumor volume and tumor cellularity varied significantly. The finding of $\leq$ ypT2N0 did not correlate with Gleason score, but did correlate with a pre-specified molecular signature, PTEN expression, and absence of cribriform/intraductal spread.

A 4-marker candidate signature (PTEN loss, nARV-7 presence, Glucocorticoid receptor (GR) high, or Ki67>10%) was predictive of resistance.

Univariate Analyses for association of pretreatment tumor characteristics with therapy effect



Diagnostic Biopsy Markers:

- ✓ **PTEN loss** enriched in persistent cancers
- ✓ **Nuclear ARV7 presence, GR high (>10%), Ki67 >10%,** correlate with persistent cancers

Marker	Pathology Stage		Fisher exact test p-value
	>ypT2N0 N(%)	$\leq$ ypT2N0 N(%)	
AR-N			
Low	13(42)	9(47)	0.77
High	18(58)	10(53)	
PTEN			
Loss	24(77)	7(37)	0.007
Intact	7(23)	12(63)	
ARV7 nuclear			
absence	19(61)	18(95)	0.009
presence	12(39)	1(5)	
GR			
Low	20(65)	17(89)	0.091
High	11(35)	2(11)	
Ki67			
$\leq$ 10%	24(77)	19(100)	0.035
>10%	7(23)	0(0)	
Clinical Stage			
cT2	6(16)	11(44)	0.02
cT3/4	32(84)	14(56)	
Biopsy			
Gleason 7	10(26)	8(32)	0.77
8-10	28(73)	17(68)	
Diagnostic			
PSA >10ng/ml	26 (68)	16 (64)	0.23

Devam Eden Neoadjuvan Çalışmaları

Name (trial number)	Location	Phase	Abbreviated Oncologic Eligibility	Treatment arms
Neoadjuvant Degarelix With or Without Apalutamide Followed by Radical Prostatectomy (ARNEO) (NCT03080116)	Leuven, Belgium	II	-Intermediate risk: at least 2 of the following factors: cT2b, biopsy GS 7, PSA 10-20ng/ml -High risk: cT≥2c and/or biopsy GS≥8 and/or PSA>20ng/ml -cN0-cN1, cM0	-Apalutamide + Degarelix -Placebo + Degarelix
Neoadjuvant Pembrolizumab Plus Androgen Axis Blockade Prior to Prostatectomy for High Risk Localized Prostate Cancer (NCT03753243)	Portland, OR, USA	II	- Any one of the following three high risk features: Gleason grade > 8-10, PSA > 20 ng/ml, cT3a -cM0	-Pembrolizumab + Enzalutamide + GNRH agonist (Single arm)
Neoadjuvant Atezolizumab-Based Combination Therapy in Men With Localized Prostate Cancer Prior to Radical Prostatectomy (NCT03821246)	San Francisco, CA, USA	II	-Only high risk patients in the safety-lead in for each cohort -Intermediate risk patients eligible once safety confirmed on interim analysis -cM0	-Atezolizumab +/- either Tocilizumab OR Etrumadenant (Non-randomized, sequential cohorts)
A Study of Neoadjuvant Hormone Therapy in Patient With Advanced Prostate Cancer Undergoing Radical Prostatectomy. (NCT03971110)	Guangzhou, China	IV	-cT3/4, cN0/1, cM0/1 (with five or fewer extra-pelvic lesions)	-Zoladex + Casodex (Single Arm)
Ibrutinib as Neoadjuvant Therapy in Localized Prostate Cancer (NCT02643667)	San Francisco, CA, USA	II	-Suitable for radical prostatectomy -cM0	-Ibrutinib (Single Arm)
Biomarkers for Neoadjuvant Pembrolizumab in Non-Metastatic Prostate Cancer Positive by 18FDG-PET Scanning (NCT04009967)	Laval, Québec, Canada	II	-Gleason Score ≥ 8, cM0 -Intraprostatic maximum standardized uptake value (SUVmax) ≥4 at 18-FDG-PET/CT exam	-Pembrolizumab (Single arm)
Neoadjuvant Hiltonol® (PolyICLC) for Prostate Cancer (NCT03262103)	New York, NY, USA	I	-Gleason 7 – 10, cT2a - cT3b adenocarcinoma of the prostate with plans for radical prostatectomy and PSA ≥ 4 ng/ml -Tumor visible on multiparametric MRI	Intratumoral injection of Poly-ICLC
177Lu-PSMA-I&T Prior to Radical Prostatectomy for Locally Advanced Disease (NCT04297410)	Petach Tikva, Israel	NA	-cT3/4 and/or Gleason score ≥8 and/or PSA ≥ 20 ng/dl -Loco-regional prostate cancer (pelvic lymphadenopathy of ≥2 cm on axial imaging) -High PSMA expression: with tracer uptake greater than normal liver (maximal SUV ≥1.5 of liver)	- 177Lu-PSMA-I&T Radionuclide (Single arm)
Neoadjuvant Therapy With Proxalutamide Combined With Androgen Deprivation Therapy (ADT) for High Risk Prostate Cancer (NCT05076851)	Nanjing, Jiangsu, China	II	- High-risk prostate cancer (cT≥2c or Gleason score ≥8 or PSA≥20ng/ml) -cM0	-Proxalutamide +ADT -Placebo + ADT

PSA Nüksü Olan Hastalarda Salvage Tedaviler

THE SALV-ENZA TRIAL (#5012)

A Phase II, Double-blind, Randomized Study of Salvage Radiation Therapy Plus Enzalutamide or Placebo for High-Risk PSA-Recurrent Prostate Cancer after Radical Prostatectomy

Phuoc T. Tran, Kathryn Lowe, Hao Wang, Hua-Ling Tsai, Daniel Y. Song, Arthur Y. Hung, Jason W.D. Hearn, Tamara Lotan, Channing Paller, Mark Markowski, Samuel Denmeade, Michael Carducci, Mario Eisenberger, Matthew Orton, Curtiland Deville, Stanley L. Liauw, Elisabeth I. Heath, Neil B. Desai, Tomasz M. Beer, Emmanuel S. Antonarakis

Salvage radiation (SRT) + ENZA monotherapy for 6 months in men with PSA-recurrent high-risk prostate cancer following prostatectomy is safe and delays PSA progression relative to SRT alone

BACKGROUND:



Does enzalutamide (ENZA) treatment without androgen deprivation therapy increase freedom-from-PSA-progression (FFPP) when combined with salvage radiation therapy (SRT) in men with recurrent prostate cancer (PCa) post-radical prostatectomy (RP)?

METHODS:

- Men with biochemically recurrent PCa after RP were randomized onto either placebo or ENZA 160 mg PO once daily for 6 months and then prostate bed radiotherapy (66.6-70.2 Gy)
- Randomization (1:1) was stratified by center, surgical margin status (R0 vs R1), PSA prior to salvage treatment (PSA ≥ 0.5 vs < 0.5 ng/mL), and pathologic Gleason sum (7 vs 8-10)
- Powered for hazard ratio (HR) 0.44 FFPP benefit with enrollment of 96 subjects
- 86 men were randomized; 43 per arm
- Median follow-up of 34 months (range 0-52)

RESULTS:

- FFPP was significantly improved with ENZA vs placebo (HR 0.42, 95% CI 0.19-0.92, $P=0.031$)
- There was differential benefit of ENZA in men with pT3 vs pT2 disease (Pinter=0.019); and R1 vs R0 (Pinter=0.023)
- Common AEs were grade 1-2 fatigue (65% ENZA vs 53% placebo) and urinary frequency (40% ENZA vs 49% placebo)




CORRESPONDENCE:

Phuoc T. Tran (phuoc.tran@som.umaryland.edu)
Emmanuel S. Antonarakis (anton401@umn.edu)

AFFILIATIONS/ENROLLING INSTITUTIONS: Johns Hopkins University, OHSU Knight Cancer Institute, University of Michigan, Wayne State University, Indiana University Health Arnett, University of Chicago, University of Texas Southwestern Medical Center, University Hospitals, UC San Francisco, University of Minnesota & University of Maryland

PSA Nüksü Olan Hastalarda Salvage Tedaviler



Sonuç

- DNA repair gen mutasyonu +, PARP inhibitörlerin kombinasyon olarak erken aşamada kullanımı
- PI3K-PTEN-AKT yolağı ilgili kombinasyon tedavileri
- PSMA ekspresyonu olanlarda Lutesyum vb +kombinasyon tedavileri daha erken aşamalarda kullanımı
- PSMA-BİTE tedavileri çoklu tedavi almış hastalarda kullanımı
- Mikrosatellit instabilite(MSI) olanlarda Checkpoint inhibitörlerin daha erken aşamada kullanımı
- Checkpoint inhibitörlerin kombinasyon(Cabozantinib vb.) olarak kastrasyona dirençli aşamada kullanımı
- Lokalize yüksek riskli ve biyokimyasal nükslerde yeni nesil AR inhibitörlerin kullanılması